

METHODS FOR TREATING HOT FLASHES AND GYNECOMASTIA**CROSS REFERENCE TO RELATED APPLICATION**

5 [0001] This application is a continuation-in-part application of United States Application Serial Number 10/609,684, filed July 1, 2003; which is a continuation-in-part application of United States Application Serial Number 10/305,363, filed November 27, 2002, and claims priority of United States Provisional Application serial number 60/333,734, filed 29 November 2001. These applications are hereby
10 incorporated in their entirety by reference herein.

FIELD OF INVENTION

[0002] This invention relates to reducing the incidence, inhibition, suppression, prevention and treatment of androgen-deprivation induced osteoporosis and/or bone fractures and/or loss of bone mineral density (BMD) in men suffering from prostate
15 cancer. More particularly, this invention relates to a method of treating, preventing, suppressing, inhibiting, or reducing the risk of developing androgen-deprivation induced osteoporosis and/or bone fractures and/or loss of BMD in men suffering from prostate cancer, comprising administering to a male subject suffering from prostate cancer an anti-estrogen agent and/or its analog, derivative, isomer,
20 metabolite, pharmaceutically acceptable salt, pharmaceutical product, hydrate, N-oxide, or any combination thereof.

BACKGROUND OF THE INVENTION

[0003] It is well established that the bone mineral density of males decrease with age.
25 Decreased amounts of bone mineral content and density correlates with decreased bone strength and predispose to fracture. The molecular mechanisms underlying the pleiotropic effects of sex-hormones in non-reproductive tissues are only beginning to be understood, but it is clear that physiologic concentrations of androgens and estrogens play an important role in maintaining bone homeostasis throughout the life-
30 cycle. Consequently, when androgen or estrogen deprivation occurs, there is a

resultant increase in the rate of bone remodeling that tilts the balance of resorption and formation in the favor of resorption, contributing to an overall loss of bone mass. In males, the natural decline in sex-hormones at maturity (direct decline in androgens as well as lower levels of estrogens derived from peripheral aromatization of androgens) is associated with the frailty of bones. This effect is also observed in males who have been castrated.

[0004] Prostate cancer is one of the most frequently diagnosed noncutaneous cancers among men in the United States. One of the approaches to the treatment of prostate cancer is by androgen deprivation. The male sex hormone, testosterone, stimulates the growth of cancerous prostatic cells and, therefore, is the primary fuel for the growth of prostate cancer. The goal of androgen deprivation is to decrease the stimulation by testosterone of the cancerous prostatic cells. Testosterone normally is produced by the testes in response to stimulation from a hormonal signal called luteinizing hormone (LH) which in turn is stimulated by luteinizing-hormone releasing hormone (LH-RH). Androgen deprivation is accomplished either surgically by bilateral orchidectomy or chemically by LH-RH agonists (LHRH \square) with or without nonsteroidal antiandrogens.

[0005] Current studies suggest that early androgen deprivation in patients with micrometastatic disease may indeed prolong survival [Messing EM, et al (1999), *N Engl J Med* 34, 1781-1788; Newling (2001), *Urology* 58(Suppl 2A), 50-55]. Moreover, androgen deprivation is being employed in numerous new clinical settings, including neoadjuvant therapy prior to radical prostatectomy, long-term adjuvant therapy for patients at high risk for recurrence following radiation or surgery, neoadjuvant therapy for radiation, and treatment of biochemical recurrence following radiation or surgery [Carroll, et al (2001), *Urology* 58, 1-4; Horwitz EM, et al (2001), *Int J Radiat Oncol Biol Phy* Mar 15;49(4), 947-56]. Thus, more prostate cancer patients have become candidates for and are being treated by androgen ablation. Moreover, these prostate cancer patients are being treated earlier and longer

than in the past, which in some cases may be as long as 10 or more years of androgen deprivation therapy.

[0006] Unfortunately, androgen deprivation therapy is fraught with significant side effects, including hot flashes, gynecomastia, osteoporosis, decreased lean muscle mass, depression and other mood changes, loss of libido, and erectile dysfunction [Stege R (2000), *Prostate Suppl* 10,38-42]. Consequently, complications of androgen blockade now contribute significantly to the morbidity, and in some cases the mortality, of men suffering from prostate cancer.

[0007] Given that more patients today are being treated by long-term androgen deprivation, osteoporosis has become a clinically important side effect in men suffering from prostate cancer undergoing androgen deprivation. Loss of bone mineral density (BMD) occurs in the majority of patients being treated by androgen deprivation by 6 months. New innovative approaches are urgently needed at both the basic science and clinical levels to decrease the incidence of androgen-deprivation induced osteoporosis in men suffering from prostate cancer.

SUMMARY OF THE INVENTION

[0008] In one embodiment, this invention provides a method of treating a subject with hot flashes, said method comprising the step of administering to said subject an anti-estrogen agent and/or its pharmaceutically acceptable salt, hydrate, N-oxide, or any combination thereof.

[0009] In one embodiment, this invention provides a method of treating a subject with gynecomastia, said method comprising the step of administering to said subject an anti-estrogen agent and/or its pharmaceutically acceptable salt, hydrate, N-oxide, or any combination thereof.

[00010] This invention relates to a method of treating androgen-deprivation induced osteoporosis in a male subject suffering from prostate cancer, the method comprising the step of administering to said subject an anti-estrogen agent and/or its analog, derivative, isomer, metabolite, pharmaceutically acceptable salt, pharmaceutical product, hydrate, N-oxide, or any combination thereof.

[00011] This invention relates to a method of preventing androgen-deprivation induced osteoporosis in a male subject suffering from prostate cancer, the method comprising the step of administering to said subject an anti-estrogen agent and/or its analog, derivative, isomer, metabolite, pharmaceutically acceptable salt, pharmaceutical product, hydrate, N-oxide, or any combination thereof.

[00012] This invention relates to a method of suppressing or inhibiting androgen-deprivation induced osteoporosis in a male subject suffering from prostate cancer, the method comprising the step of administering to said subject an anti-estrogen agent and/or its analog, derivative, isomer, metabolite, pharmaceutically acceptable salt, pharmaceutical product, hydrate, N-oxide, or any combination thereof.

[00013] This invention relates to a method of reducing the risk of developing androgen-deprivation induced osteoporosis in a male subject suffering from prostate cancer, the method comprising the step of administering to said subject an anti-estrogen agent and/or its analog, derivative, isomer, metabolite, pharmaceutically acceptable salt, pharmaceutical product, hydrate, N-oxide, or any combination thereof.

[00014] This invention relates to a method of treating androgen-deprivation induced loss of BMD in a male subject suffering from prostate cancer, the method comprising the step of administering to said subject an anti-estrogen agent and/or its analog, derivative, isomer, metabolite, pharmaceutically acceptable salt, pharmaceutical product, hydrate, N-oxide, or any combination thereof.

[00015] This invention relates to a method of preventing androgen-deprivation induced loss of BMD in a male subject suffering from prostate cancer, the method comprising the step of administering to said subject an anti-estrogen agent and/or its
5 analog, derivative, isomer, metabolite, pharmaceutically acceptable salt, pharmaceutical product, hydrate, N-oxide, or any combination thereof.

[00016] This invention relates to a method of suppressing or inhibiting androgen-deprivation induced loss of BMD in a male subject suffering from prostate cancer,
10 the method comprising the step of administering to said subject an anti-estrogen agent and/or its analog, derivative, isomer, metabolite, pharmaceutically acceptable salt, pharmaceutical product, hydrate, N-oxide, or any combination thereof.

[00017] This invention relates to a method of reducing the risk of developing
15 androgen-deprivation induced loss of BMD in a male subject suffering from prostate cancer, the method comprising the step of administering to said subject an anti-estrogen agent and/or its analog, derivative, isomer, metabolite, pharmaceutically acceptable salt, pharmaceutical product, hydrate, N-oxide, or any combination thereof.

[00018] This invention relates to a method of treating androgen-deprivation induced bone fractures in a male subject suffering from prostate cancer, the method comprising the step of administering to said subject an anti-estrogen agent and/or its
20 analog, derivative, isomer, metabolite, pharmaceutically acceptable salt,
25 pharmaceutical product, hydrate, N-oxide, or any combination thereof.

[00019] This invention relates to a method of preventing androgen-deprivation induced bone fractures in a male subject suffering from prostate cancer, the method comprising the step of administering to said subject an anti-estrogen agent and/or its
30 analog, derivative, isomer, metabolite, pharmaceutically acceptable salt,

pharmaceutical product, hydrate, N-oxide, or any combination thereof.

[00020] This invention relates to a method of suppressing or inhibiting androgen-deprivation induced bone fractures in a male subject suffering from prostate cancer, the method comprising the step of administering to said subject an anti-estrogen agent and/or its analog, derivative, isomer, metabolite, pharmaceutically acceptable salt, pharmaceutical product, hydrate, N-oxide, or any combination thereof.

[00021] This invention relates to a method of reducing the risk of developing androgen-deprivation induced bone fractures in a male subject suffering from prostate cancer, the method comprising the step of administering to said subject an anti-estrogen agent and/or its analog, derivative, isomer, metabolite, pharmaceutically acceptable salt, pharmaceutical product, hydrate, N-oxide, or any combination thereof.

[00022] In one embodiment the anti-estrogen is a selective estrogen receptor modulator (SERM). In another embodiment, the anti-estrogen is a triphenylethylene. In another embodiment, the anti-estrogen is Toremifene.

[00023] In one embodiment, the anti-estrogen is administered at a daily dosage of about 20 mg. In another embodiment, the anti-estrogen is administered at a daily dosage of about 40 mg. In another embodiment, the anti-estrogen is administered at a daily dosage of about 60 mg. In another embodiment, the anti-estrogen is administered at a daily dosage of about 80 mg.

[00024] In one embodiment, the anti-estrogen is Toremifene, which is administered at a daily dosage of about 20 mg. In another embodiment, the anti-estrogen is Toremifene, which is administered at a daily dosage of about 40 mg. In another embodiment, the anti-estrogen is Toremifene, which is administered at a daily dosage of about 60 mg. In another embodiment, the anti-estrogen is

Toremifene, which is administered at a daily dosage of about 80 mg.

[00025] The present invention provides a safe and effective method for treating, preventing, suppressing, inhibiting or reducing the risk of developing androgen-deprivation induced osteoporosis and/or loss of BMD and is particularly useful for treating male subjects suffering from prostate cancer having an elevated risk of developing androgen-deprivation induced osteoporosis and/or loss of BMD.

10

BRIEF DESCRIPTION OF THE FIGURES

[00026] The present invention will be understood and appreciated more fully from the following detailed description taken in conjunction with the appended figures which depict:

15 **Figure 1:** Effect of Toremifene on C-telopeptide of rat collagen I (RatLaps ELISA).

Figure 2: Effect of Toremifene on serum osteocalcin levels. A) 10 and 30 days; B) 60 and 120 days.

20

DETAILED DESCRIPTION OF THE INVENTION

[00027] This invention provides: 1) a method of treating a subject with hot flashes; 2) a method of treating a subject with gynecomastia; 3) a method of treating androgen-deprivation induced osteoporosis in a male subject suffering from prostate cancer; 4) a method of preventing androgen-deprivation induced osteoporosis in a male subject suffering from prostate cancer; 5) a method of suppressing or inhibiting androgen-deprivation induced osteoporosis in a male subject suffering from prostate cancer; 6) a method of reducing the risk of developing androgen-deprivation induced osteoporosis in a male subject suffering from prostate cancer; 7) a method of treating androgen-deprivation induced loss of BMD in a male subject suffering from

30

prostate cancer; 8) a method of preventing androgen-deprivation induced loss of BMD in a male subject suffering from prostate cancer; 9) a method of suppressing or inhibiting androgen-deprivation induced loss of BMD in a male subject suffering from prostate cancer; 10) a method of reducing the risk of developing androgen-deprivation induced loss of BMD in a male subject suffering from prostate cancer; 11) a method of treating androgen-deprivation induced bone fractures in a male subject suffering from prostate cancer; 12) a method of preventing androgen-deprivation induced bone fractures in a male subject suffering from prostate cancer; 13) a method of suppressing or inhibiting androgen-deprivation induced bone fractures in a male subject suffering from prostate cancer; 14) a method of reducing the risk of developing androgen-deprivation induced bone fractures in a male subject suffering from prostate cancer by administering to the subject an anti-estrogen agent and/or its analog, derivative, isomer, metabolite, pharmaceutically acceptable salt, pharmaceutical product, hydrate, N-oxide, or any combination thereof.

15

[00028] As provided herein, the results demonstrate that administration of an anti-estrogen, such as, for example, Toremifene, at a daily dosage of from about 20 mg to about 80 mg, is bone sparing. This is determined by measuring the levels of bone-specific serum markers that indicate bone resorption and formation. Further, this invention demonstrates that an anti-estrogen, such as, for example, Toremifene (and/or 17 -Estradiol), increases bone mineral density.

20

[00029] Toremifene is an example of a triphenylalkylene compound described in US. Patent Nos. 4,696,949 and 5,491,173 to Toivola et al., the disclosures of which are incorporated herein by reference. The parenteral and topical administration to mammalian subjects of formulations containing Toremifene is described in U.S. Patent No. 5,571,534 to Jalonen et al. and in U.S. Patent No. 5,605,700 to DeGregorio et al., the disclosures of which are incorporated herein by reference.

25

[00030] In one embodiment the anti-estrogen that treats, prevents, suppresses, inhibits

30

or reduces the risk of developing androgen-deprivation induced osteoporosis and/or loss of BMD is a selective estrogen receptor modulator (SERM) and/or its analog, derivative, isomer, metabolite, pharmaceutically acceptable salt, pharmaceutical product, hydrate, N-oxide, or any combination thereof.

5

[00031] In one embodiment, the SERMs that are encompassed by the present invention include, but are not limited to the following embodiments:

triphenylalkylenes such as triphenylethylenes, which include Tamoxifen, Droloxifene, Toremifene, Idoxifene, Clomiphene, Enclomiphene and Zuclomiphene; benzothiphenes such as Raloxifene and LY 353381; benzopyran derivatives such as EM 800 (SCH 57050) and its metabolite EM 652; naphthalene derivatives such as Lasofoxifene (CP 336,156); chromans such as Levormeloxifene or their analogs, derivatives, isomers, or metabolites thereof, or their pharmaceutically acceptable salts, esters, N-oxides, or mixtures thereof.

15

[00032] As contemplated herein, other embodiments of anti-estrogens that are encompassed by the present invention include but are in no way limited to the following embodiments: Cycladiene, Merck Index, 10th ed. #3085 and U.S. Pat. No. 2,464,203 and U.S. Pat. No. 2,465,505; Nafoxidine, USAN and USP Dictionary of Drug Names, p. 327 (1983); CI-680, Unlisted Drugs, 28(10): 169(o) (1976); CI-628, Unlisted Drugs, 26(7): 106(l) (1974); CN-55,945-27, or nitromifene citrate, Unlisted Drugs, 27(12): 194(n) (1975); R2323 or 13-ethyl-17a-ethynl- 17B-hydroxygona-4,9,11-trien-3-one, Unlisted Drugs, 23(3): 34(b) (1971); MER-25; U-11,555A; U-11,100A; ICI-46,669 and ICI-46,474; all shown in L. Terenius, et al., "Aspects on the Mode of Action of Antiestrogens and Antiprogestrogens," Hormones and Antagonists. Gynec. Invest. 3: 98; Diphenol hydrochrysene; erythro-MEA; and Park Davis CN-55,945; all disclosed in C. Geynet, et al., "Estrogens and Antiestrogens," Hormones and Antagonists. Gynec. Invest. 3: 12-13 (1972); Allenolic acid and cyclofenyl, disclosed in C. Geynet, et al., Hormones and Antagonists. Gynec. Invest. 3: 17 (1972); Chlorotrianisene, Merck Index, 10th ed., #2149; Ethamoxytriphetol,

30

Merck Index, 10th ed., #3668; and Triparanol, Merck Index, 10th ed., #9541 and U.S. Pat. No. 2,914,562.

5 [00033] Various embodiments of dosage ranges are contemplated by this invention. In one embodiment, the dosage is in the range of 1-80 mg/day. In another embodiment, the dosage is in the range of 5-80 mg/day. In another embodiment the dosage is in the range of 35-66 mg/day. In another embodiment the dosage is in the range of 20-80 mg/day. In another embodiment the dosage is in the range of 20-60 mg/day. In another embodiment the dosage is in the range of 40-60 mg/day. In another
10 embodiment the dosage is in a range of 45-60 mg/day. In another embodiment the dosage is in the range of 15-25 mg/day. In another embodiment the dosage is in the range of 55-65 mg/day. In one embodiment, the dosage is 20 mg/day. In another embodiment, the dosage is 40 mg/day. In another embodiment, the dosage is 60 mg/day. In another embodiment, the dosage is 80 mg/day.

15

[00034] Accordingly, the present invention provides a method of treating androgen-deprivation induced osteoporosis in a male subject suffering from prostate cancer, the method comprising the step of administering to said subject an anti-estrogen agent and/or its analog, derivative, isomer, metabolite, pharmaceutically acceptable salt,
20 pharmaceutical product, hydrate, N-oxide, or any combination thereof, at a dosage of about 20 mg to about 80 mg per day of the anti-estrogen.

[00035] In another embodiment, the present invention provides a method of preventing androgen-deprivation induced osteoporosis in a male subject suffering
25 from prostate cancer, the method comprising the step of administering to said subject an anti-estrogen agent and/or its analog, derivative, isomer, metabolite, pharmaceutically acceptable salt, pharmaceutical product, hydrate, N-oxide, or any combination thereof, at a dosage of about 20 mg to about 80 mg per day of the anti-estrogen.

30

[00036] In another embodiment, the present invention provides a method of suppressing or inhibiting androgen-deprivation induced osteoporosis in a male subject suffering from prostate cancer, the method comprising the step of administering to said subject an anti-estrogen agent and/or its analog, derivative, isomer, metabolite, pharmaceutically acceptable salt, pharmaceutical product, hydrate, N-oxide, or any combination thereof, at a dosage of about 20 mg to about 80 mg per day of the anti-estrogen.

[00037] In another embodiment, the present invention provides a method of reducing the risk of developing androgen-deprivation induced osteoporosis in a male subject suffering from prostate cancer, the method comprising the step of administering to said subject an anti-estrogen agent and/or its analog, derivative, isomer, metabolite, pharmaceutically acceptable salt, pharmaceutical product, hydrate, N-oxide, or any combination thereof, at a dosage of about 20 mg to about 80 mg per day of the anti-estrogen.

[00038] In another embodiment, the present invention provides a method of treating androgen-deprivation induced loss of BMD in a male subject suffering from prostate cancer, the method comprising the step of administering to said subject an anti-estrogen agent and/or its analog, derivative, isomer, metabolite, pharmaceutically acceptable salt, pharmaceutical product, hydrate, N-oxide, or any combination thereof, at a dosage of about 20 mg to about 80 mg per day of the anti-estrogen.

[00039] In another embodiment, the present invention provides a method of preventing androgen-deprivation induced loss of BMD in a male subject suffering from prostate cancer, the method comprising the step of administering to said subject an anti-estrogen agent and/or its analog, derivative, isomer, metabolite, pharmaceutically acceptable salt, pharmaceutical product, hydrate, N-oxide, or any combination thereof, at a dosage of about 20 mg to about 80 mg per day of the anti-estrogen.

[00040] In another embodiment, the present invention provides a method of suppressing or inhibiting androgen-deprivation induced loss of BMD in a male subject suffering from prostate cancer, the method comprising the step of administering to said subject an anti-estrogen agent and/or its analog, derivative, isomer, metabolite, pharmaceutically acceptable salt, pharmaceutical product, hydrate, N-oxide, or any combination thereof, at a dosage of about 20 mg to about 80 mg per day of the anti-estrogen.

10 [00041] In another embodiment, the present invention provides a method of reducing the risk of developing androgen-deprivation induced loss of BMD in a male subject suffering from prostate cancer, the method comprising the step of administering to said subject an anti-estrogen agent and/or its analog, derivative, isomer, metabolite, pharmaceutically acceptable salt, pharmaceutical product, hydrate, N-oxide, or any
15 combination thereof, at a dosage of about 20 mg to about 80 mg per day of the anti-estrogen.

[00042] In another embodiment, the present invention provides a method of treating androgen-deprivation induced bone fractures in a male subject suffering from prostate cancer, the method comprising the step of administering to said subject an
20 anti-estrogen agent and/or its analog, derivative, isomer, metabolite, pharmaceutically acceptable salt, pharmaceutical product, hydrate, N-oxide, or any combination thereof, at a dosage of about 20 mg to about 80 mg per day of the anti-estrogen.

25 [00043] In another embodiment, the present invention provides a method of preventing androgen-deprivation induced bone fractures in a male subject suffering from prostate cancer, the method comprising the step of administering to said subject an anti-estrogen agent and/or its analog, derivative, isomer, metabolite, pharmaceutically acceptable salt, pharmaceutical product, hydrate, N-oxide, or any
30 combination thereof, at a dosage of about 20 mg to about 80 mg per day of the anti-

estrogen.

[00044] In another embodiment, the present invention provides a method of suppressing or inhibiting androgen-deprivation induced bone fractures in a male subject suffering from prostate cancer, the method comprising the step of administering to said subject an anti-estrogen agent and/or its analog, derivative, isomer, metabolite, pharmaceutically acceptable salt, pharmaceutical product, hydrate, N-oxide, or any combination thereof, at a dosage of about 20 mg to about 80 mg per day of the anti-estrogen.

10

[00045] In another embodiment, the present invention provides a method of reducing the risk of developing androgen-deprivation induced bone fractures in a male subject suffering from prostate cancer, the method comprising the step of administering to said subject an anti-estrogen agent and/or its analog, derivative, isomer, metabolite, pharmaceutically acceptable salt, pharmaceutical product, hydrate, N-oxide, or any combination thereof, at a dosage of about 20 mg to about 80 mg per day of the anti-estrogen.

15

[00046] In one embodiment the anti-estrogen is a selective estrogen receptor modulator (SERM). In another embodiment, the anti-estrogen is a triphenylethylene. In another embodiment, the anti-estrogen is Toremifene.

20

[00047] In one embodiment, the anti-estrogen is administered at a daily dosage of about 20 mg. In another embodiment, the anti-estrogen is administered at a daily dosage of about 40 mg. In another embodiment, the anti-estrogen is administered at a daily dosage of about 60 mg. In another embodiment, the anti-estrogen is administered at a daily dosage of about 80 mg.

25

[00048] In one embodiment, the anti-estrogen is Toremifene, which is administered at a daily dosage of about 20 mg. In another embodiment, the anti-

30

estrogen is Toremifene, which is administered at a daily dosage of about 40 mg. In another embodiment, the anti-estrogen is Toremifene, which is administered at a daily dosage of about 60 mg. In another embodiment, the anti-estrogen is Toremifene, which is administered at a daily dosage of about 80 mg.

5

[00049] Osteoporosis is a systemic skeletal disease, characterized by low bone mass and deterioration of bone tissue, with a consequent increase in bone fragility and susceptibility to fracture. In osteoporotic patients, bone strength is abnormal, with a resulting increase in the risk of fracture. Osteoporosis depletes both the calcium and the protein collagen normally found in the bone, resulting in either abnormal bone quality or decreased bone density. Bones that are affected by osteoporosis can fracture with only a minor fall or injury that normally would not cause a bone fracture. The fracture can be either in the form of cracking (as in a hip fracture) or collapsing (as in a compression fracture of the spine). The spine, hips, and wrists are common areas of osteoporosis bone fractures, although fractures can also occur in other skeletal areas.

10

15

20

[00050] BMD is a measured calculation of the true mass of bone. The absolute amount of bone as measured by bone mineral density (BMD) generally correlates with bone strength and its ability to bear weight. By measuring BMD, it is possible to predict fracture risk in the same manner that measuring blood pressure can help predict the risk of stroke.

25

30

[00051] BMD in one embodiment can be measured by known bone-mineral content mapping techniques. Bone density of the hip, spine, wrist, or calcaneus may be measured by a variety of techniques. The preferred method of BMD measurement is dual-energy x-ray densitometry (DXA). BMD of the hip, antero-posterior (AP) spine, lateral spine, and wrist can be measured using this technology. Measurement at any site predicts overall risk of fracture, but information from a specific site is the best predictor of fracture at that site. Quantitative computerized tomography (QCT)

is also used to measure BMD of the spine. See for example, "Nuclear Medicine: Quantitative Procedures". by Wahner H W, Dunn W L, Thorsen H C, et al, published by Toronto Little, Brown & Co., 1983, (see pages 107-132). An article entitled "Assessment of Bone Mineral Part 1" appeared in the Journal of Nuclear
5 Medicine, pp 1134-1141, (1984). Another article entitled "Bone Mineral Density of The Radius" appeared in Vol. 26, No. 11, (1985) Nov. Journal of Nuclear Medicine at pp 13-39. Abstracts on the use of gamma cameras for bone-mineral content measurements are (a) S. Hoory et al, Radiology, Vol. 157(P), p. 87 (1985), and (b) C.
R. Wilson et al, Radiology, Vol. 157(P), p. 88 (1985).

10

[00052] The present invention provides a safe and effective method for treating, preventing, suppressing, inhibiting or reducing the risk of developing androgen-deprivation induced osteoporosis and/or loss of BMD and is particularly useful for treating male subjects suffering from prostate cancer having an elevated risk of
15 developing androgen-deprivation induced osteoporosis. In one embodiment, the male subject is a mammalian subject. In another embodiment, the male subject is a human subject.

20

[00053] Furthermore, the anti-estrogens presented herein are effective at treating, suppressing or inhibiting osteopenia accompanied by bone loss. "Osteopenia" refers to decreased calcification or density of bone. This is a term which encompasses all skeletal systems in which such a condition is noted.

25

[00054] As contemplated herein, the present invention relates to the use of an anti-estrogen compound and/or its analog, derivative, isomer, metabolite, pharmaceutically acceptable salt, pharmaceutical product, hydrate, N-oxide, or combinations thereof for treating, preventing, suppressing, inhibiting or reducing the risk of developing androgen-deprivation induced osteoporosis and/or loss of BMD at a dosage of about 20 mg to about 80 mg per day of the anti-estrogen. Thus, in one
30 embodiment, the methods of the present invention comprise administering an analog

of the anti-estrogen. In another embodiment, the methods of the present invention comprise administering a derivative of the anti-estrogen. In another embodiment, the methods of the present invention comprise administering an isomer of the anti-estrogen. In another embodiment, the methods of the present invention comprise administering a metabolite of the anti-estrogen. In another embodiment, the methods of the present invention comprise administering a pharmaceutically acceptable salt of the anti-estrogen. In another embodiment, the methods of the present invention comprise administering a pharmaceutical product of the anti-estrogen. In another embodiment, the methods of the present invention comprise administering a hydrate of the anti-estrogen. In another embodiment, the methods of the present invention comprise administering an N-oxide of the anti-estrogen. In another embodiment, the methods of the present invention comprise administering any of a combination of an analog, derivative, isomer, metabolite, pharmaceutically acceptable salt, pharmaceutical product, hydrate or N-oxide of the anti-estrogen.

[00055] As defined herein, the term "isomer" includes, but is not limited to, optical isomers and analogs, structural isomers and analogs, conformational isomers and analogs, and the like.

[00056] In one embodiment, this invention encompasses the use of various optical isomers of the anti-estrogen compound. It will be appreciated by those skilled in the art that the anti-estrogens of the present invention contain at least one chiral center. Accordingly, the anti-estrogens used in the methods of the present invention may exist in, and be isolated in, optically-active or racemic forms. Some compounds may also exhibit polymorphism. It is to be understood that the present invention encompasses any racemic, optically-active, polymorphic, or stereoisomeric form, or mixtures thereof, which form possesses properties useful in the treatment of androgen-related conditions described herein. In one embodiment, the anti-estrogens are the pure (R)-isomers. In another embodiment, the anti-estrogens are the pure (S)-isomers. In another embodiment, the anti-estrogens are a mixture of the (R) and the

(S) isomers. In another embodiment, the anti-estrogens are a racemic mixture comprising an equal amount of the (R) and the (S) isomers. It is well known in the art how to prepare optically-active forms (for example, by resolution of the racemic form by recrystallization techniques, by synthesis from optically-active starting materials, 5 by chiral synthesis, or by chromatographic separation using a chiral stationary phase).

[00057] The invention includes “pharmaceutically acceptable salts” of amino-substituted compounds with organic and inorganic acids, for example, citric acid and hydrochloric acid. The invention also includes N-oxides of the amino substituents of 10 the compounds described herein. Pharmaceutically acceptable salts can also be prepared from the phenolic compounds by treatment with inorganic bases, for example, sodium hydroxide. Also, esters of the phenolic compounds can be made with aliphatic and aromatic carboxylic acids, for example, acetic acid and benzoic acid esters.

15 [00058] This invention further includes derivatives of the anti-estrogens. The term “derivatives” includes but is not limited to ether derivatives, acid derivatives, amide derivatives, ester derivatives and the like. In addition, this invention further includes hydrates of the anti-estrogen compounds. The term “hydrate” includes but is not 20 limited to hemihydrate, monohydrate, dihydrate, trihydrate and the like.

[00059] This invention further includes metabolites of the anti-estrogen compounds. The term “metabolite” means any substance produced from another substance by metabolism or a metabolic process.

25 [00060] This invention further includes pharmaceutical products of the anti-estrogen compounds. The term “pharmaceutical product” means a composition suitable for pharmaceutical use (pharmaceutical composition), as defined herein.

30 [00061] In addition, the invention encompasses pure (Z)- and (E)- isomers of the anti-

estrogen compounds defined herein and mixtures thereof as well as pure (RR,SS)- and (RS,SR)-enantiomer couples and mixtures thereof.

Pharmaceutical Compositions

5 [00062] In one embodiment, the methods of the present invention comprise administering a pharmaceutical composition comprising the anti-estrogen and/or its analog, derivative, isomer, metabolite, pharmaceutically acceptable salt, pharmaceutical product, hydrate, N-oxide, or any combination thereof; and a pharmaceutically acceptable carrier. The pharmaceutical composition is administered
10 to a male subject suffering from prostate cancer; for treating and/or preventing androgen-deprivation induced osteoporosis and/or loss of BMD; for suppressing or inhibiting androgen-deprivation induced osteoporosis and/or loss of BMD; and/or for reducing the risk of developing androgen-deprivation induced osteoporosis and/or loss of BMD in the male subject.

15 [00063] As used herein, "pharmaceutical composition" means a "therapeutically effective amount" of the active ingredient, i.e. the anti-estrogen, together with a pharmaceutically acceptable carrier or diluent. A "therapeutically effective amount" as used herein refers to that amount which provides a therapeutic effect for a given
20 condition and administration regimen.

[00064] The pharmaceutical compositions containing the anti-estrogen can be administered to a subject by any method known to a person skilled in the art, such as parenterally, paracancerally, transmucosally, transdermally, intramuscularly,
25 intravenously, intradermally, subcutaneously, intraperitoneally, intraventricularly, intracranially, intravaginally or intratumorally.

[00065] In one embodiment, the pharmaceutical compositions are administered orally, and are thus formulated in a form suitable for oral administration, i.e. as a
30 solid or a liquid preparation. Suitable solid oral formulations include tablets,

capsules, pills, granules, pellets and the like. Suitable liquid oral formulations include solutions, suspensions, dispersions, emulsions, oils and the like. In one embodiment of the present invention, the anti-estrogen compounds are formulated in a capsule. In accordance with this embodiment, the compositions of the present invention comprise, in addition to the anti-estrogen active compound and the inert carrier or diluent, a hard gelating capsule.

[00066] Further, in another embodiment, the pharmaceutical compositions are administered by intravenous, intraarterial, or intramuscular injection of a liquid preparation. Suitable liquid formulations include solutions, suspensions, dispersions, emulsions, oils and the like. In one embodiment, the pharmaceutical compositions are administered intravenously, and are thus formulated in a form suitable for intravenous administration. In another embodiment, the pharmaceutical compositions are administered intraarterially, and are thus formulated in a form suitable for intraarterial administration. In another embodiment, the pharmaceutical compositions are administered intramuscularly, and are thus formulated in a form suitable for intramuscular administration.

[00067] Further, in another embodiment, the pharmaceutical compositions are administered topically to body surfaces, and are thus formulated in a form suitable for topical administration. Suitable topical formulations include gels, ointments, creams, lotions, drops and the like. For topical administration, the anti-estrogen agents or their physiologically tolerated derivatives such as salts, esters, N-oxides, and the like are prepared and applied as solutions, suspensions, or emulsions in a physiologically acceptable diluent with or without a pharmaceutical carrier.

[00068] Further, in another embodiment, the pharmaceutical compositions are administered as a suppository, for example a rectal suppository or a urethral suppository. Further, in another embodiment, the pharmaceutical compositions are

administered by subcutaneous implantation of a pellet. In a further embodiment, the pellet provides for controlled release of anti-estrogen agent over a period of time.

5 [00069] In another embodiment, the active compound can be delivered in a vesicle, in particular a liposome (see Langer, Science 249:1527-1533 (1990); Treat et al., in Liposomes in the Therapy of Infectious Disease and Cancer, Lopez- Berestein and Fidler (eds.), Liss, New York, pp. 353-365 (1989); Lopez-Berestein, *ibid.*, pp. 317-327; see generally *ibid.*).

10 [00070] As used herein "pharmaceutically acceptable carriers or diluents" are well known to those skilled in the art. The carrier or diluent may be a solid carrier or diluent for solid formulations, a liquid carrier or diluent for liquid formulations, or mixtures thereof.

15 [00071] Solid carriers/diluents include, but are not limited to, a gum, a starch (e.g. corn starch, pregeletanized starch), a sugar (e.g., lactose, mannitol, sucrose, dextrose), a cellulosic material (e.g. microcrystalline cellulose), an acrylate (e.g. polymethylacrylate), calcium carbonate, magnesium oxide, talc, or mixtures thereof.

20 [00072] For liquid formulations, pharmaceutically acceptable carriers may be aqueous or non-aqueous solutions, suspensions, emulsions or oils. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, and injectable organic esters such as ethyl oleate. Aqueous carriers include water, alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media. Examples of oils are
25 those of petroleum, animal, vegetable, or synthetic origin, for example, peanut oil, soybean oil, mineral oil, olive oil, sunflower oil, and fish-liver oil.

[00073] Parenteral vehicles (for subcutaneous, intravenous, intraarterial, or intramuscular injection) include sodium chloride solution, Ringer's dextrose, dextrose
30 and sodium chloride, lactated Ringer's and fixed oils. Intravenous vehicles include

fluid and nutrient replenishers, electrolyte replenishers such as those based on Ringer's dextrose, and the like. Examples are sterile liquids such as water and oils, with or without the addition of a surfactant and other pharmaceutically acceptable adjuvants. In general, water, saline, aqueous dextrose and related sugar solutions, and glycols such as propylene glycols or polyethylene glycol are preferred liquid carriers, particularly for injectable solutions. Examples of oils are those of petroleum, animal, vegetable, or synthetic origin, for example, peanut oil, soybean oil, mineral oil, olive oil, sunflower oil, and fish-liver oil.

[00074] In addition, the compositions may further comprise binders (e.g. acacia, cornstarch, gelatin, carbomer, ethyl cellulose, guar gum, hydroxypropyl cellulose, hydroxypropyl methyl cellulose, povidone), disintegrating agents (e.g. cornstarch, potato starch, alginic acid, silicon dioxide, croscarmellose sodium, crospovidone, guar gum, sodium starch glycolate), buffers (e.g., Tris-HCl, acetate, phosphate) of various pH and ionic strength, additives such as albumin or gelatin to prevent absorption to surfaces, detergents (e.g., Tween 20, Tween 80, Pluronic F68, bile acid salts), protease inhibitors, surfactants (e.g. sodium lauryl sulfate), permeation enhancers, solubilizing agents (e.g., glycerol, polyethylene glycerol), anti-oxidants (e.g., ascorbic acid, sodium metabisulfite, butylated hydroxyanisole), stabilizers (e.g. hydroxypropyl cellulose, hydroxypropylmethyl cellulose), viscosity increasing agents (e.g. carbomer, colloidal silicon dioxide, ethyl cellulose, guar gum), sweeteners (e.g. aspartame, citric acid), preservatives (e.g., Thimerosal, benzyl alcohol, parabens), lubricants (e.g. stearic acid, magnesium stearate, polyethylene glycol, sodium lauryl sulfate), flow-aids (e.g. colloidal silicon dioxide), plasticizers (e.g. diethyl phthalate, triethyl citrate), emulsifiers (e.g. carbomer, hydroxypropyl cellulose, sodium lauryl sulfate), polymer coatings (e.g., poloxamers or poloxamines), coating and film forming agents (e.g. ethyl cellulose, acrylates, polymethacrylates) and/or adjuvants.

[00075] In one embodiment, the pharmaceutical compositions provided herein are controlled-release compositions, i.e. compositions in which the anti-estrogen

compound is released over a period of time after administration. Controlled- or sustained-release compositions include formulation in lipophilic depots (e.g. fatty acids, waxes, oils). In another embodiment, the composition is an immediate-release composition, i.e. a composition in which all of the anti-estrogen compound is released immediately after administration.

[00076] In yet another embodiment, the pharmaceutical composition can be delivered in a controlled release system. For example, the agent may be administered using intravenous infusion, an implantable osmotic pump, a transdermal patch, liposomes, or other modes of administration. In one embodiment, a pump may be used (see Langer, supra; Sefton, CRC Crit. Ref. Biomed. Eng. 14:201 (1987); Buchwald et al., Surgery 88:507 (1980); Saudek et al., N. Engl. J. Med. 321:574 (1989). In another embodiment, polymeric materials can be used. In yet another embodiment, a controlled release system can be placed in proximity to the therapeutic target, i.e., the brain, thus requiring only a fraction of the systemic dose (see, e.g., Goodson, in Medical Applications of Controlled Release, supra, vol. 2, pp. 115-138 (1984). Other controlled-release systems are discussed in the review by Langer (Science 249:1527-1533 (1990).

[00077] The compositions may also include incorporation of the active material into or onto particulate preparations of polymeric compounds such as polylactic acid, polyglycolic acid, hydrogels, etc, or onto liposomes, microemulsions, micelles, unilamellar or multilamellar vesicles, erythrocyte ghosts, or spheroplasts.) Such compositions will influence the physical state, solubility, stability, rate of *in vivo* release, and rate of *in vivo* clearance.

[00078] Also comprehended by the invention are particulate compositions coated with polymers (e.g. poloxamers or poloxamines) and the compound coupled to antibodies directed against tissue-specific receptors, ligands or antigens or coupled to ligands of tissue-specific receptors.

[00079] Also comprehended by the invention are compounds modified by the covalent attachment of water-soluble polymers such as polyethylene glycol, copolymers of polyethylene glycol and polypropylene glycol, carboxymethyl cellulose, dextran, polyvinyl alcohol, polyvinylpyrrolidone or polyproline. The modified compounds are known to exhibit substantially longer half-lives in blood following intravenous injection than do the corresponding unmodified compounds (Abuchowski et al., 1981; Newmark et al., 1982; and Katre et al., 1987). Such modifications may also increase the compound's solubility in aqueous solution, eliminate aggregation, enhance the physical and chemical stability of the compound, and greatly reduce the immunogenicity and reactivity of the compound. As a result, the desired *in vivo* biological activity may be achieved by the administration of such polymer-compound adducts less frequently or in lower doses than with the unmodified compound.

[00080] The preparation of pharmaceutical compositions which contain an active component is well understood in the art, for example by mixing, granulating, or tablet-forming processes. The active therapeutic ingredient is often mixed with excipients which are pharmaceutically acceptable and compatible with the active ingredient. For oral administration, the anti-estrogen agents or their physiologically tolerated derivatives such as salts, esters, N-oxides, and the like are mixed with additives customary for this purpose, such as vehicles, stabilizers, or inert diluents, and converted by customary methods into suitable forms for administration, such as tablets, coated tablets, hard or soft gelatin capsules, aqueous, alcoholic or oily solutions. For parenteral administration, the anti-estrogen agents or their physiologically tolerated derivatives such as salts, esters, N-oxides, and the like are converted into a solution, suspension, or emulsion, if desired with the substances customary and suitable for this purpose, for example, solubilizers or other.

[00081] An active component can be formulated into the composition as neutralized pharmaceutically acceptable salt forms. Pharmaceutically acceptable salts include the acid addition salts (formed with the free amino groups of the polypeptide or antibody molecule), which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like. Salts formed from the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, 2-ethylamino ethanol, histidine, procaine, and the like.

10

[00082] For use in medicine, the salts of the anti-estrogens are pharmaceutically acceptable salts. Other salts may, however, be useful in the preparation of the compounds according to the invention or of their pharmaceutically acceptable salts. Suitable pharmaceutically acceptable salts of the compounds of this invention include acid addition salts which may, for example, be formed by mixing a solution of the compound according to the invention with a solution of a pharmaceutically acceptable acid such as hydrochloric acid, sulphuric acid, methanesulphonic acid, fumaric acid, maleic acid, succinic acid, acetic acid, benzoic acid, oxalic acid, citric acid, tartaric acid, carbonic acid or phosphoric acid.

20

[00083] As defined herein, "contacting" means that the anti-estrogen compound of the present invention is introduced into a sample containing the enzyme in a test tube, flask, tissue culture, chip, array, plate, microplate, capillary, or the like, and incubated at a temperature and time sufficient to permit binding of the anti-estrogen to the enzyme. Methods for contacting the samples with the anti-estrogen or other specific binding components are known to those skilled in the art and may be selected depending on the type of assay protocol to be run. Incubation methods are also standard and are known to those skilled in the art.

25

[00084] In another embodiment, the term “contacting” means that the anti-estrogen compound of the present invention is introduced into a subject receiving treatment, and the anti-estrogen compound is allowed to come in contact with the androgen receptor *in vivo*.

5

[00085] As used herein, the term “treating” includes preventative as well as disorder remitative treatment. As used herein, the terms “reducing”, “suppressing” and “inhibiting” have their commonly understood meaning of lessening or decreasing. As used herein, the term “progression” means increasing in scope or severity, advancing, growing or becoming worse. As used herein, the term “recurrence” means the return of a disease after a remission.

10

[00086] As used herein, the term “administering” refers to bringing a subject in contact with an anti-estrogen compound of the present invention. As used herein, administration can be accomplished *in vitro*, i.e. in a test tube, or *in vivo*, i.e. in cells or tissues of living organisms, for example humans. In one embodiment, the present invention encompasses administering the compounds of the present invention to a subject.

15

[00087] In one embodiment, the methods of the present invention comprise administering an anti-estrogen compound as the sole active ingredient. However, also encompassed within the scope of the present invention are methods for hormone therapy, for treating prostate cancer, for delaying the progression of prostate cancer, and for preventing and/or treating the recurrence of prostate cancer, which comprise administering the anti-estrogen compounds in combination with one or more therapeutic agents. These agents include, but are not limited to: LHRH analogs, reversible antiandrogens (such as bicalutamide or flutamide), additional anti-estrogens, anticancer drugs, 5-alpha reductase inhibitors, aromatase inhibitors, progestins, selective androgen receptor modulators (SARMS) or agents acting through other nuclear hormone receptors.

20

25

30

[00088] Thus, in one embodiment, the methods of the present invention include using compositions and pharmaceutical compositions comprising an anti-estrogen, in combination with an LHRH analog. In another embodiment, the methods of the present invention include using compositions and pharmaceutical compositions comprising an anti-estrogen, in combination with a reversible antiandrogen. In another embodiment, the methods of the present invention include using compositions and pharmaceutical compositions comprising an anti-estrogen, in combination with an additional anti-estrogen. In another embodiment, the methods of the present invention include using compositions and pharmaceutical compositions comprising an anti-estrogen, in combination with an anticancer drug. In another embodiment, the methods of the present invention include using compositions and pharmaceutical compositions comprising an anti-estrogen, in combination with a 5-alpha reductase inhibitor. In another embodiment, the methods of the present invention include using compositions and pharmaceutical compositions comprising an anti-estrogen, in combination with an aromatase inhibitor. In another embodiment, the methods of the present invention include using compositions and pharmaceutical compositions comprising an anti-estrogen, in combination with a progestin. In another embodiment, the methods of the present invention include using compositions and pharmaceutical compositions comprising an anti-estrogen, in combination with a SARM. In another embodiment, the methods of the present invention include using compositions and pharmaceutical compositions comprising an anti-estrogen, in combination with an agent acting through other nuclear hormone receptors.

[00089] The following examples are presented in order to more fully illustrate the preferred embodiments of the invention. They should in no way, however, be construed as limiting the broad scope of the invention.

EXPERIMENTAL DETAILS SECTION

EXAMPLE 1

Effect of Toremifene on bone turnover in human male subjects

5 [00090] In a Phase IIa clinical trial to determine whether Toremifene has chemopreventive activity against prostate cancer, 18 men with high-grade prostatic intraepithelial neoplasia (HGPIN) were treated with 60mg/d of Toremifene for 4 months. At Day 120 there was a significant reduction from baseline in serum calcium (mean -0.12, p=0.005) and at both day 60 and day 120, alkaline phosphatase was
10 significantly decreased compared to baseline (mean=-18.7 at Day 60 and -21.0 at Day 120, and p<0.001 for both visits).

[00091] These clinical data demonstrate that the anti-estrogen Toremifene showed estrogenic effects on bone favorably affecting bone turnover markers in men.

15

EXAMPLE 2: EFFECT OF TOREMIFENE ON BONE IN MALE RATS

Drug Delivery System

[00092] The test article, positive control and placebo were delivered by ALZA pumps
20 manufactured by Durect Corporation (Cupertino, CA). Pumps were implanted in a subcutaneous pocket using appropriate surgical technique. The pumps employed in this study deliver a continuous rate of drug over a 30-day period with Toremifene formulated to release 1.8 mg/day (2 mL pump) and 17- μ g/day. -Estradiol (positive control) is released at 70 μ g/day. Data provided by the manufacturer of the pumps
25 validates the constant rate of drug delivery over a 28 day period, and suggests that the constant rate can be expected for several additional days. Animals were anesthetized and pump replacement was performed for each dosage group on days 31, 61, and 91 to provide drug administration over a 120 day period. Every animal on study had a pellet implanted to control for potential confounding variables
30 associated with surgery for implantation.

Study Groups:

[00093] Adult male Sprague-Dawley rats (14-weeks old at start of study), weighing approximately 0.35 kg, were used in the experiments. This study employed five test groups of 12 rats. Treatment groups represent placebo control (castrated and sham operated), 17- β -estradiol (castrated) and Toremifene (castrated and sham operated). 5 animals from each treatment group were sacrificed at day 60 and day 120 and bone metabolism markers were measured and bone were harvested for biomechanical strength and density testing.

10

Table 1. Treatment groups for Effects of Toremifene on Bone

Group	Drug	Sham	Castrated	# of animals	Sacrifice Schedule (days)
1	Placebo control	✓		5	60
2	Placebo control	✓		5	120
3	Toremifene (1.75 mg/day)	✓		5	60
4	Toremifene (1.75 mg/day)	✓		5	120
5	Estradiol (0.07 mg/day)		✓	5	60
6	Estradiol (0.07 mg/day)		✓	5	120
7	Placebo control		✓	5	60
8	Placebo control		✓	5	120
9	Toremifene (1.75 mg/day)		✓	5	60

10	Toremifene (1.75 mg/day)		✓	5	120
----	-----------------------------	--	---	---	-----

Observations:

- 5 [00094] Clinical Observations: Cage-side observations were made and recorded once weekly (daily cage checks note any exceptions). Effects on vital functions were observed. Any animals found moribund or presenting with signs of acute toxicity were anesthetized by an intramuscular ketamine/xylazine (87/13) injection and euthanized by abdominal aorta exsanguinations.
- 10 [00095] Body Weight: Body weights were taken in triplicate and averaged once per week.
- 15 [00096] Sacrifices were made at 60 and 120-days. Each sacrifice involved 5 rats from each treatment group, and 2 rats at 30, 60, 90, and 120 days to verify drug in plasma (refer to Table 1 for group allocation to treatment and sacrifice). Animals were anesthetized by an injection of ketamine/xylazine (87/13) and sacrificed by abdominal aorta exsanguinations. Blood collected at exsanguinations was processed to collect serum.
- 20 [00097] Bone remodeling was assessed per serum markers to analyze bone resorption and bone formation of treatment rats. The tests listed in Table 2 indicate the bone remodeling category and the amounts of serum required for analysis. Serum chemistry analysis of blood calcium, phosphorus, bilirubin and creatinine levels, and bone-specific alkaline phosphatase, were performed at AniLytics, Inc. (Gaithersburg, MD).
- 25

Table 2. Assays for serum markers of bone remodeling in mouse

Remodeling Category	Tests Run	Test Supplier	Minimum Serum Required
Bone Formation	Rat-MID Osteocalcin ELISA	OSTEOMET ER	20ul
	Chemistry Analysis of Calcium, Phosphorus, Billirubin, Creatine	AniLytics, Inc.	70ul
	Bone Spec. Phosphatase	AniLytics, Inc.	50ul
Bone Resorption	Serum CrossLaps One Step ELISA	OSTEOMET ER	20ul
	(Deoxy)Pyridinolin e Crosslink in serum	METRA	25ul
	RatTRAP ELISA (tartrate-resistant Acid P)	SBA Sciences	50ul

5 [00098] Specimen collection included femoral bones in addition to blood. Blood was
 processed for serum. Serum was aliquoted and frozen at -80°C until analysis. Serum
 tests were performed at GTx, Inc. for osteocalcin levels (bone formation marker) and
 c-telopeptide (bone resorption marker). After the femur was removed from each
 animal, it was stripped of extraneous tissue, and stored at -20°C until biomechanical
 10 strength and bone mineral density analysis.

RANDOMIZATION/ASSIGNMENT TO GROUPS:

[00099] Prior to study, 10 groups were randomized to treatments, then sixty (60) 12 week-old rats were ordered, ear-tagged and weighed by the Manager of Animal Resources. Sixty animals were then randomized to the 10 treatment groups of 6 animals each. An ANOVA was performed to establish the presence or absence of significant difference in body weight (within 10% of average body weight is acceptable). The population was already restricted and could not be restricted further. If there was a significant difference, it was duly noted, but changes in assignment were not made. Bone parameters were not normalized to body weight. Values for bone parameters are reported as absolute and evaluated based on group means compared to the appropriate controls.

COMPOUNDS

[000100] Test Article 1:

15 Identification: Toremifene

Description: White crystalline powder, that is extracted from Toremifene tablets as active ingredient.

[000101] Test Article 2 (Positive Control):

20 Identification: 17- β -Estradiol (Sigma Aldrich).

EXAMPLE 3**EFFECT OF TOREMIFENE ON BONE DENSITY AND SERUM**

25 **MARKERS FOR BONE REMODELING IN MALE RATS**

[000102] The purpose of this study is to determine whether administration of Toremifene to mature male rats is bone sparing as can presently be measured by the levels of bone-specific serum markers that indicate bone resorption and formation (where 17- β -Estradiol is used as a positive control). The effect of Toremifene (and

30

17-□ -Estradiol) on androgen deprivation-induced bone loss was also determined through bone density and mechanical strength testing.

[000103] It is well established that the bone mineral density of males, similar to that of
5 females, decreases with age. Decreased amounts of bone mineral content and density correlates with decreased bone strength, and predisposes to fracture. The molecular mechanism underlying the pleiotropic effects of sex-hormones in non-reproductive tissues are only beginning to be understood, but it is clear that physiologic concentrations of androgens and estrogens play an important role in maintaining bone
10 homeostasis throughout the life-cycle. Consequently, when androgen and/or estrogen deprivation occurs, there is a resultant increase in the rate of bone remodeling that tilts the balance of resorption and formation in the favor of resorption, contributing to the overall loss of bone mass. In males, the natural decline in sex-hormone at maturity (direct decline in androgens as well as lower levels of estrogens derived
15 from peripheral aromatization of androgens) is associated with the frailty of bones. This effect is also observed in males who have been castrated. Previous studies demonstrate in mice that pharmacological therapy with selective estrogen receptor modulator (SERM) compounds in androgen deficient males has a positive effect on bone mineral density and that bone can be maintained or even increased (Broulik,
20 *Horm Metab Res*, 2000; 32: 181-184). It is suggested that in humans, this intervention would halt the onset of osteoporosis or at least reduce its severity. Importantly, recent human data indicates that estrogen loss in elderly males is more critical for short-term bone changes than loss of androgens. It is believed that the selective estrogen receptor modulator Toremifene (trade name is AcapodeneTM and
25 generic name is toremifene) will have similar bone-building properties when compared with other nonsteroidal anti-estrogens (e.g. Tamoxifene, (Broulik, *Horm Metab Res*, 2000; 32: 181-184), and will be a clinically useful drug for preventing osteoporosis and maintaining bone density in aging males. The model used herein is an orchidectomy model, which is an experimental model used to mimic the type of
30 androgen deprivation that would be caused by, for example, LHRH agonist therapy in

prostate cancer.

MATERIALS AND METHODS

Study Design

5 [000104] Male Sprague-Dawley rats (Harlan Sprague Dawley) were placed on study at
14-weeks of age. They were randomized and divided into five treatment groups:
vehicle only (placebo, or P) after sham operation, vehicle only after orchidectomy
(Orx), Toremifene (5 mg/kg/day) after sham-operation, and Toremifene after
orchidectomy, and 17- β -estradiol (0.5 mg/kg/day) after orchidectomy. Test articles
10 were delivered subcutaneously by Alzet pumps. Pumps were re-implanted every 30
days until the end of the study. Five to six animals from each group were sacrificed
after 60 and 120 days of treatment, and femurs were harvested, stripped from soft
tissue and muscle, and stored individually in polypropylene vials at -80°C until
analysis of bone density and mechanical strength testing. Additionally, serum was
15 collected on days 15, 30, 60, 90 and 120, to measure markers of bone metabolism and
to perform serum chemistry.

[000105] After collection, serum was divided into 3 aliquots and stored at -80°C until
analysis. 125 μl for serum biochemistry performed by AniLytics, Gaithersburg, MD
20 (Bone-Specific Alkaline Phosphatase, Calcium, Phosphorus, Creatinine and
Bilirubin). 100 μl for serum Rat Osteocalcin ELISA and Serum RatLaps ELISA for
C-terminal telopeptides determination. The rest of the collected serum was left for
repeated test if necessary.

25 [000106] Bones were stripped from soft tissue and muscles and stored individually in
15 ml vials at -80°C , until further testing.

30

Analysis of Bone Turnover Markers from Rat Serum**5 Serum RatLaps ELISA, Osteometer BioTech A/S, Denmark**

[000107] The assay was performed in duplicates, each 20 μ l of serum as well as standards and control. Briefly, 100 μ l of Biotinylated RatLaps Antigen was incubated 30 min in each well, strips were washed and 20 μ l of Standard, Control and unknown samples were added to appropriate wells, followed by 100 μ l of
10 primary antibody (polyclonal Ab raised against a synthetic peptide EKSQDGGR specific for a part of the C-terminal telopeptide α 1 chain of rat type I collagen). After overnight incubation at 4 C, the wells were washed and 100 μ l of Peroxidase conjugated Goat anti-Rabbit IgG Antibody was added to each well and incubated 60 min. After washing the strips 100 μ l of the Chromogenic Substrate Solution was
15 added to each well, and after 15 min the reaction was stopped and the absorbance was measured at 450 nm. The means of the duplicate absorbance determination were calculated, and a standard curve was constructed using 4-parametric logistic curve fit. Sample RatLaps concentrations were determined by interpolation.

20 Rat Osteocalcin EIA, Biomedical Technologies, MA

[000108] This assay is specific for rat Osteocalcin and both carboxylated and decarboxylated rat Osteocalcin are recognized. A monoclonal antibody against the N-terminal region of Osteocalcin is bound to the polystyrene wells. The assay was performed in duplicates. 100 μ l of standards, control and 1 to 20 diluted samples
25 were added to appropriate wells and incubated overnight at 4 C. After washing the strips, 100 μ l of Goat polyclonal antibody of high specificity for C-terminus of rat Osteocalcin was added and incubated 60 min at 37 C. After washing, 100 μ l of Donkey anti-goat IgG peroxidase conjugated antibody was added and incubated 60 min at 22 C. The wells were washed and a mix of TMB and peroxide solution was

added and incubated 30 min at 22 C in the dark. The reaction was stopped, and the absorbance was measured at 450 nm. A standard curve was constructed using 4-parametric logistic curve fit. Sample osteocalcin concentrations were determined by interpolation.

5

Bone Analysis Methodology

Peripheral Quantitative Computed Tomography (pQCT)

[000109] Right femurs were subjected to pQCT analysis using a Stratec XCT-RM with associated software (Stratec Medizintechnik GmbH, Pforzheim, Germany. Software version 5.40 C). Femurs were scanned at two sites, 20% and 50% of the total femoral length measured from the distal femur. The position was verified using scout views, and scan results from two 0.5 mm slices perpendicular to the long axis of the femur shaft were recorded.

15 Mechanical Testing

[000110] Using a Material Testing System (Model 5501R Instron Corp., Canton, MA), two types of mechanical testing were performed on the right femur. The load and extension curves were collected by the machine software (Merlin II, Instron Corp.). All tests were conducted using a 5 kN load cell at a constant loading rate of 6 mm/min. The general applications of the tests were as described in Turner and Burr (*Bone*, 1993; 14: 595-608) and Ke et al (*Bone*, 1998; 23, 249-255).

20

Compression Test of the Distal Femur

[000111] A compression test was used to determine the mechanical properties of the distal femur. The distal femur test specimen was obtained by removing a 3 mm segment directly proximal to the distal condyle using a slow speed diamond saw with constant saline irrigation. An electronic caliper was used to measure the average

25

anterior/posterior diameter (a), medial/lateral diameter (b), and height (h) of the bone. The extrinsic parameters, maximal load (F_u), stiffness (S) and energy (W), were obtained from the load and extension curve. The following intrinsic parameters were calculated from the measured values: Cross sectional area (CSA) = $(\pi * a * b) / 4$;
 5 Ultimate Strength (σ) = F_u / CSA ; Elastic modulus (E) = $S / (CSA / h)$; Toughness (T) = $W / (CSA * h)$.

RESULTS/DISCUSSION

Analysis of Serum Markers of Bone Turnover

10 [000112] Bone turnover markers have been demonstrated as an effective, validated tool for the clinical scientist to monitor bone activity. Ultimately, the data of greatest consequence to evaluate novel therapeutics aimed at osteoporosis treatment and prevention is a demonstrated improvement in the quality of bone itself. However, because changes in bone turnover markers correlate well with bone strength testing,
 15 in the present study C-telopeptide and osteocalcin levels were analyzed for interim analysis and supplementary data to support of the effectiveness of treatment.

C-telopeptide of Type I collagen:

[000113] As demonstrated in Figure 1, C-telopeptide levels in orchidectomized animals
 20 were slightly increased by 22 and 9.5% over the placebo 15 and 30 day groups, respectively, indicating that, after castration, bone resorptive activity is increased and type I collagen is being degraded, with fragments containing the cross-linked molecules released into the blood. Further, treatment of orchidectomized animals with toremifene and the positive control, 17- β -estradiol, reduced the C-telopeptide
 25 levels to at or below control values (i.e. unorchidectomized, placebo-treated animals), with the 30 day treatment trending to significance for the 17- β -estradiol group.

[000114] Because toremifene and 17- β -estradiol decrease the level of C-telopeptide in the serum, the data indicate that these agents are acting to prevent the resorption of
 30 bone induced by androgen deprivation.

Osteocalcin:

[000115] Similar to other published literature, Applicants found that osteocalcin levels were increased by castration. Toremifene significantly reduced the osteocalcin levels in castrated animals to intact control levels ($p < 0.05$ at 15 days, and $p < 0.02$ at 30 days, Figure 2A). The increase in osteocalcin levels were most pronounced 15 days after castration, although Toremifene and 17- β -estradiol continued to significantly reduce osteocalcin levels to below that of intact control rats for up to 120 days (Figure 2B). These results indicate that the rate of bone formation in males is upregulated following orchidectomy to compensate for increased resorptive activity. 17- β -Estradiol, and the selective estrogen receptor modulator toremifene, stabilize the bone resorption and formation processes, hence decreasing overall osteocalcin levels that are detectable in serum.

15 Biomechanical Analysis of Bone

[000116] Androgen deficiency, induced by castration, has been used as a model of male osteoporosis. In this model, most of the bone loss occurs in cancellous bone. To further understand the effects of toremifene on bone mineral density and mechanical strength, bones were harvested upon completion of the in-life study phase and sent to SkeleTech (Bothell, WA) for testing. All bones were thawed in physiological saline prior to analysis. Statistical analysis was performed using SAS software (SAS Institute, Cary, NC). One-way analysis of variance (group) was performed. Individual group differences were ascertained with Dunnett's procedure using treatment group 2 (Castrated + Placebo) as the reference group. A p value of < 0.05 was considered significant. Where appropriate, a p value of < 0.1 is noted as a trend (when the treatment results are in the direction of the positive control).

Distal Right Femur Cancellous pQCT:

[000117] Total bone mineral content and density were lower, though not statistically

significant, in Orx animals when compared to Sham-operated animals. Both Toremifene and 17- β -Estradiol treatment appeared to reduce total bone area and increase total bone mineral density. Cancellous bone mineral content and density in Orx animals were 34% less than Sham-operated animals. Toremifene partially prevented this decrease whereas 17- β -Estradiol was able to fully prevent this cancellous bone loss. None of the parameters produced statistically significant differences from the Orx group due to the small sample size and the large variations in the measured results. The results are summarized in Table 3.

10 **Table 3: Summary of the Distal Right Femur Cancellous pQCT**

Treatm ent Group		Total Bone Content (mg/mm)	Total Bone Area (mm ²)	Total Bone Density (mg/cm ³)	Trabecu lar Content (mg/mm)	Trabecu lar Area (mm ²)	Trabecu lar Density (mg/cm ³)
Sham (n = 6)		11.47	17.41	658.27	1.65	7.83	209.72
	S. E. M. .	0.89	0.97	32.81	0.43	0.44	52.17
	p< 0.0 5	n.s.	n.s	n.s	n.s	n.s	n.s
Cx (n = 5)	M e a n	11.07	17.38	639.50	1.08	7.82	138.38
	S. E. M. .	0.50	1.26	47.07	0.22	0.57	31.88
	p < 0. 0 5	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.

Cx + Toremifene (n = 5)	M e a n	10.69	15.96	669.74	1.13	7.18	157.56
	S. E. M .	0.48	0.67	21.95	0.22	0.30	32.77
	p < 0. 0 5	n.s	n.s	n.s	n.s	n.s	n.s
Cx + EST (n = 5)	M e a n	11.01	15.95	690.74	1.52	7.17	210.28
	S. E. M .	0.76	0.98	26.68	0.50	0.44	62.69
	p < 0. 0 5	n.s	n.s	n.s	n.s	n.s	n.s

p<0.05 vs. TG2.

n.s. = not significant

n.a. = not applicable

5 **Compression Test of the Distal Femur:**

[000118] Mechanical strength was tested at the distal femur, a site rich in cancellous bone. Maximum load, stiffness, ultimate strength, and elastic modulus were lower in Orx animals when compared to Sham-operated animals. Toremifene treatment of Orx animals improved various parameters to a level that was better than Sham-operated animals. 17- β -Estradiol treatment showed statistically significant improvements in these parameter values. Cross-sectional area was decreased in toremifene and 17- β -Estradiol -treated Orx animals as expected from pQCT data. The results are summarized in Table 3.

Table 4: Summary of the Distal Right Femur Compression Test

Treatment Group		Maximum Load	Stiffness	Energy	Ultimate Strength	Elastic Modulus	Toughness	CSA	Height
		(N)	(N/mm)	(mJ)	(N/mm ²)	(MPa)	(MJ/mm ³)	(mm ²)	(mm)
Sham (n = 6)		420.04	3400.22	43.10	18.95	481.38	0.62	22.32	3.13
	S.D.	123.75	944.50	13.09	5.86	152.38	0.18	1.64	0.14
	p < 0.05	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.
Cx (n = 5)		386.80	2538.96	49.97	16.73	341.02	0.68	23.19	3.14
	S.D.	70.43	887.56	18.45	3.33	109.52	0.23	1.28	0.10
	p < 0.05	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.
Cx + Toremi fene (n = 5)		467.01	3856.71	56.96	22.13	579.07	0.86	21.30	3.15
	S.D.	125.89	1303.56	17.07	6.34	211.70	0.30	1.43	0.15
	p < 0.05	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.
Cx + EST (n = 5)		613.61	4842.67	58.28	31.23	753.44	0.98	19.75	3.06
	S.D.	72.10	740.18	18.39	4.76	136.97	0.34	1.14	0.12

	p < 0.0 5	*	*	n.s.	*	*	n.s.	*	n.s.
--	-----------------	---	---	------	---	---	------	---	------

p<0.05 vs. TG2.

n.s. = not significant

n.a. = not applicable

5 SUMMARY

[000119] Androgen deficiency model resulted in animals that had increased levels of the bone resorption marker C-telopeptide as well as osteocalcin in serum. Treatment with toremifene and 17- β -estradiol significantly reduced the levels of these serum markers following castration. Further, androgen deficiency resulted in a 34% loss of cancellous bone mineral content and density. Importantly, the use of toremifene partially prevented this loss. As predicted, estrogen was very effective in preventing cancellous bone loss due to androgen deficiency. Additionally, the compression test of the distal femur showed improved strength parameters in orchidectomized groups treated with toremifene, and statistically significant improvements with estrogen treatment. These measurements are partially correlated with total bone mineral density at that site. In conclusion, the data presented herein suggests that the selective estrogen receptor modulator toremifene would have a positive effect for bone improvement in men undergoing androgen-deprivation therapy for prostate cancer.

[000120] It will be appreciated by a person skilled in the art that the present invention is not limited by what has been particularly shown and described hereinabove. Rather, the scope of the invention is defined by the claims which follow: